

NutriCam

Gilt
Rearer 2



Klas: Volledige Varkgroeivoer / Class: Complete Pig Grower Feed

V30182

| Voedingstof | Eenheid / Unit | | Nutrient |
|--------------------|----------------|------|-------------------|
| Proteïen (Min) | 140 | g/kg | Protein (Min) |
| Lisien (Min) | 7.0 | g/kg | Lysine (Min) |
| Metionien (Min) | 2.5 | g/kg | Methionine (Min) |
| Vog (Maks) | 120 | g/kg | Moisture (Max) |
| Vet (Min) | 25 | g/kg | Fat (Min) |
| Vesel (Maks) | 80 | g/kg | Fibre (Max) |
| Kalsium (Min/Maks) | 6.0-10.0 | g/kg | Calcium (Min/Max) |
| Fosfor (Min) | 5.0 | g/kg | Phosphorus (Min) |
| Sout (Maks) | 5.0 | g/kg | Salt (Max) |

PRODUKTESKRYWING / PRODUCT DESCRIPTION

'n Volledig volvoer wat ontwikkel is om in die voedingsbehoefes van groeiende vervangingsoggies te voorsien. Hierdie voer verseker goeie skeletontwikkeling en ideale vetreserwes.

A complete gilt rearing feed developed to meet the nutrient requirements of growing replacement gilts. This feed supports good skeletal development and ideal fat reserves.

VOERINSTRUKSIES / FEEDING INSTRUCTIONS

Voer *ad lib* vanaf 168 tot 203 - 210 dae ouderdom. Die liggaamsgewig by 203 tot 210 dae moet 136 tot 145 kg wees wanneer die sog vir die tweede keer op hitte kom. Hierdie voer moet as deel van 'n voedingsprogram vir soggies wees en saam met NutriCam Gilt Rearer 1 gevoer word. Voer 0,5 – 1,5 kg van die PorCare Maxi-Flush S-diet saam met die NutriCam Gilt Rearer 2 vir 5 tot 7 dae voor dekking totdat die 'spoeleffek' bereik word. Soggies moet weekliks tydens die grootmaak geweeg word om seker te maak dat hulle die groeikurve volg. Om optimale groei te verseker moet daar voldoende voederspasie per vark wees. Voeders moet altyd skoon gehou word en vars water moet ten alle tye beskikbaar wees.

Feed *ad lib* from 168 to 203 - 210 days of age. Target body weight at 203 to 210 days should be 136 to 145 kg at the second heat. This feed should be used as part of a phased gilt rearing programme with the Nutricam Gilt Rearer 1. Feed 0,5 to 1,5 kg of the PorCare Maxi-Flush S diet on top of the NutriCam Gilt Rearer 2 for 5 to 7 days before mating to achieve the 'flushing effect'. Gilts should be weighed weekly during rearing to be sure that they are following growth curve. For optimum growth there must be sufficient feeder space per pig. Feeders must always be kept clean and water must always be available.

AANBEVOLE INNAMES / RECOMMENDED INTAKE

120 kg per soggie.

120 kg per gilt.